



<https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assesment-hcp.html>

Key Points

- In general, asymptomatic HCP who have had a higher-risk exposure do not require work restriction, regardless of vaccination status, if they do not develop symptoms or test positive for SARS-CoV-2.

Test-based strategy

HCP who are symptomatic could return to work after the following criteria are met:

- Resolution of fever without the use of fever-reducing medications, **and**
- Improvement in symptoms (e.g., cough, shortness of breath), **and**
- Results are negative from at least two consecutive respiratory specimens collected 48 hours apart (total of two negative specimens) tested using an antigen test or NAAT.

HCP who are not symptomatic could return to work after the following criteria are met:

- Results are negative from at least two consecutive respiratory specimens collected 48 hours apart (total of two negative specimens) tested using an antigen test or NAAT.

Return to Work Criteria for HCP with SARS-CoV-2 Infection

The following are criteria to determine when HCP with SARS-CoV-2 infection could return to work and are influenced by severity of symptoms and presence of immunocompromising conditions. After returning to work, HCP should self-monitor for symptoms and seek re-evaluation from occupational health if symptoms recur or worsen. If symptoms recur (e.g., rebound) these HCP should be restricted from work and follow recommended practices to prevent transmission to others (e.g., use of well-fitting source control) until they again meet the healthcare criteria below to return to work unless an alternative diagnosis is identified.

HCP with **mild to moderate illness** who are **not moderately to severely immunocompromised** could return to work after the following criteria have been met:

- At least 7 days have passed *since symptoms first appeared* if a negative viral test* is obtained within 48 hours prior to returning to work (or 10 days if testing is not performed or if a positive test at day 5-7), **and**
- At least 24 hours have passed *since last fever* without the use of fever-reducing medications, **and**
- Symptoms (e.g., cough, shortness of breath) have improved.

*Either a NAAT (molecular) or antigen test may be used. If using an antigen test, HCP should have a negative test obtained on day 5 and again 48 hours later

HCP who were asymptomatic throughout their infection and are **not moderately to severely immunocompromised** could return to work after the following criteria have been met:

- At least 7 days have passed since the date of their first positive viral test if a negative viral test* is obtained within 48 hours prior to returning to work (or 10 days if testing is not performed or if a positive test at day 5-7).

*Either a NAAT (molecular) or antigen test may be used. If using an antigen test, HCP should have a negative test obtained on day 5 and again 48 hours later