


Take home point: Among older women residing in nursing homes, administration of cranberry capsules, compared with placebo, resulted in no significant difference in presence of bacteriuria plus pyuria over 1 year (1)

Cranberry products have been shown to prevent adherence of P-fimbriated E. coli to uroepithelial cells. Not all UTIs are E. coli.

No statistically significant difference between Cranberry group and Placebo group (2).



UTI Facts

- Bacteriuria is prevalent in 25-50% of female nursing home residents and pyuria is present in 90% of those with bacteriuria (1)
- Bacteriuria is prevalent in 25-50% of female nursing home residents and pyuria is present in 90% of those with bacteriuria (3).



- A randomized trial of antibiotic treatment versus no treatment of bacteriuria in nursing home women showed no decrease in genitourinary morbidity or mortality with treatment (4).
- As many as 75% of prescriptions for UTI in LTCF residents are given to individuals who do not meet criteria for UTI (5).

Resources

- 1. <u>Infectious Diseases in Older Adults of Long-term Care Facilities: Update on Approach</u> <u>to Diagnosis and Management – PMC (nih.gov)</u>
- 2. <u>Effect of Cranberry Capsules on Bacteriuria Plus Pyuria among Older Women in</u> <u>Nursing Homes: A Randomized Clinical Trial – PMC (nih.gov)</u>
- 3. <u>Tests for Urinary Tract Infection in Nursing Home Residents | Urology | JAMA | JAMA Network</u>
- 4. <u>Prospective randomized comparison of therapy and no therapy for asymptomatic</u> <u>bacteriuria in institutionalized elderly women – PubMed (nih.gov)</u>
- 5. <u>Challenges Assessing Nursing Home Residents with Advanced Dementia for Suspected</u> <u>Urinary Tract Infections – PMC (nih.gov)</u>
- 6. FDA Announces Qualified Health Claim for Certain Cranberry Products and Urinary Tract Infections | FDA