



## Efficacy of Oral Risperidone, Haloperidol, or Placebo for Symptoms of Delirium

Among Patients in Palliative Care: A Randomized Clinical Trial | End of Life,

### Abstract

Hospice, Palliative Care | JAMA Internal Medicine | JAMA Network

**Importance** Antipsychotics are widely used for distressing symptoms of delirium, but efficacy has not been established in placebo-controlled trials in palliative care.

**Objective** To determine efficacy of risperidone or haloperidol relative to placebo in relieving target symptoms of delirium associated with distress among patients receiving palliative care.

**Design, Setting, and Participants** A double-blind, parallel-arm, dose-titrated randomized clinical trial was conducted at 11 Australian inpatient hospice or hospital palliative care services between August 13, 2008, and April 2, 2014, among participants with life-limiting illness, delirium, and a delirium symptoms score (sum of Nursing Delirium Screening Scale behavioral, communication, and perceptual items) of 1 or more.

**Interventions** Age-adjusted titrated doses of oral risperidone, haloperidol, or placebo solution were administered every 12 hours for 72 hours, based on symptoms of delirium. Patients also received supportive care, individualized treatment of delirium precipitants, and subcutaneous midazolam hydrochloride as required for severe distress or safety.

**Main Outcome and Measures** Improvement in mean group difference of delirium symptom score (severity range, 0-6) between baseline and day 3. Five a priori secondary outcomes: delirium severity, midazolam use, extrapyramidal effects, sedation, and survival.

**Results** Two hundred forty-seven participants (mean [SD] age, 74.9 [9.8] years; 85 women [34.4%]; 218 with cancer [88.3%]) were included in intention-to-treat analysis (82 receiving risperidone, 81 receiving haloperidol, and 84 receiving placebo). In the primary intention-to-treat analysis, participants in the risperidone arm had delirium symptom scores that were significantly higher than those among participants in the placebo arm (on average 0.48 Units higher; 95% CI, 0.09-0.86;  $P = .02$ ) at study end. Similarly, for those in the haloperidol arm, delirium symptom scores were on average 0.24 Units higher (95% CI, 0.06-0.42;  $P = .009$ ) than in the placebo arm. Compared with placebo, patients in both active arms had more extrapyramidal effects (risperidone, 0.73; 95% CI, 0.09-1.37;  $P = .03$ ; and haloperidol, 0.79; 95% CI, 0.17-1.41;  $P = .01$ ). Participants in the placebo group had better overall survival than those receiving haloperidol (hazard ratio, 1.73; 95% CI, 1.20-2.50;  $P = .003$ ), but this was not significant for placebo vs risperidone (hazard ratio, 1.29; 95% CI, 0.91-1.84;  $P = .14$ ).

**Conclusions and Relevance** In patients receiving palliative care, individualized management of delirium precipitants and supportive strategies result in lower scores and shorter duration of target distressing delirium symptoms than when risperidone or haloperidol are added.



Efficacy of Oral Risperidone, Haloperidol, or Placebo for Symptoms of Delirium

Among Patients in Palliative Care: A Randomized Clinical Trial | End of Life,

<https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2588810>  
Hospice, Palliative Care | JAMA Internal Medicine | JAMA Network