

Randomized Study of Basal-Bolus Insulin Therapy in the Inpatient Management of Patients

With Type 2 Diabetes (RABBIT 2 Trial) | Diabetes Care | American Diabetes Association

(diabetesjournals.org)

• **CONCLUSIONS**—Treatment with insulin glargine and glulisine resulted in significant improvement in glycemic control compared with that achieved with the use of SSI alone. Our study indicates that a basal-bolus insulin regimen is preferred over SSI in the management of non-critically ill, hospitalized patients with type 2 diabetes.

Randomized Study of Basal-Bolus Insulin Therapy in the Inpatient Management of Patients

With Type 2 Diabetes Undergoing General Surgery (RABBIT 2 Surgery) | Diabetes Care |

American Diabetes Association (diabetesjournals.org)

CONCLUSIONS

Basal-bolus treatment with glargine once daily plus glulisine before meals improved
glycemic control and reduced hospital complications compared with SSI in general
surgery patients. Our study indicates that a basal-bolus insulin regimen is preferred
over SSI in the hospital management of general surgery patients with type 2 diabetes.

 $\frac{https://shmpublications.onlinelibrary.wiley.com/pb-assets/assets/15535606/jhmbf01319-sup-0001-1648471892367.ppt}{0001-1648471892367.ppt}$

Sliding-Scale-Insulin-in-hospitalDownload

The latest on inpatient glucose control | ACP Hospitalist (acponline.org)