



thickened liquids does not lower aspiration pneumonia risk



Why This is Unnecessary and Potentially Harmful

- Thickened liquids are associated with higher rates of dehydration, fever, and urinary tract infections in comparison to thin liquids.⁷
- Perceived QOL of patients using thickened liquids is significantly lower.⁸
- Compliance with thickened liquids is low (~35% after 5 days).⁹
- No significant difference in the risk of developing pneumonia in patients with access to free water when compared to those with thickened liquids.^{2,5,6,8,10}

Journal of
Hospital Medicine

@JHospMedicine | #TWDFNR

shm.

References

Based on Lippert WC, Chadha R, Sweigart JR. The Use of Thickened Liquids in Treating Hospitalized Adult Patients with Dysphagia. *J Hosp Med.* 2019 May;**14**(5):315-317.

- 1.Foley N, Teasell R, Salter K, Kruger E, Martino R. Dysphagia treatment post stroke: a systematic review of randomised controlled trials. *Age Ageing.* 2008;**37**(3):258-264.
- 2.Loeb MB, Becker M, Eady A, Walker-Dilks C. Interventions to prevent aspiration pneumonia in older adults: a systematic review. *J Am Geriatr Soc.* 2003;**51**(7):1018-1022.
- 3.Steele CM, Alsanei WA, Ayanikalath S, et al. The influence of food texture and liquid consistency modification on swallowing physiology and function: a systematic review. *Dysphagia.* 2015;**30**(1):2-26.
- 4.Clave P, de Kraa M, Arreola V, et al. The effect of bolus viscosity on swallowing function in neurogenic dysphagia. *Aliment Pharmacol Ther.* 2006;**24**(9):1385-1394.
- 5.Gillman A, Winkler R, Taylor NF. Implementing the free water protocol does not result in aspiration pneumonia in carefully selected patients with dysphagia: a systematic review. *Dysphagia.* 2017;**32**(3):345-361.
- 6.Kaneoka A, Pisegna JM, Saito H, et al. A systematic review and meta-analysis of pneumonia associated with thin liquid vs. thickened liquid intake in patients who aspirate. *Clin Rehabil.* 2017;**31**(8):1116-1125.
- 7.Robbins J, Gensler G, Hind J, et al. Comparison of 2 interventions for liquid aspiration on pneumonia incidence: a randomized trial. *Ann Intern Med.* 2008;**148**(7):509-518.
- 8.Carlaw C, Finlayson H, Beggs K, et al. Outcomes of a pilot water protocol project in a rehabilitation setting. *Dysphagia.* 2012;**27**(3):297-306.
- 9.Leiter AE WJ. Compliance of geriatric dysphagic patients with safe-swallowing instructions. *J Med Speech Lang Pathol.* 1996;**4**(4):289-300.
- 10.DePippo KL, Holas MA, Reding MJ, Mandel FS, Lesser ML. Dysphagia therapy following stroke: a controlled trial. *Neurology.* 1994;**44**(9):1655-1660.
- 11.Ashford J, McCabe D, Wheeler-Hegland K, et al. Evidence-based systematic review: Oropharyngeal dysphagia behavioral treatments. Part III—impact of dysphagia treatments on populations with neurological disorders. *J Rehabil Res Dev.* 2009;**46**(2):195-204.

<https://shmpublications.onlinelibrary.wiley.com/pb-assets/assets/15535606/jhmbf01367-sup-0001-164841>

[8140057.ppt](#)