



## Prepare for Survey!! The focus will likely be on Infection Control, Life Safety, and Emergency Preparedness.

The **Office of the Inspector General** issued a report about Colorado nursing homes in February 2024. Full OIG report can be accessed at the link below. I encourage you to read this report paying special attention to the Findings section; there are pictures.

***“The Findings detailed in our report show that nursing homes in Colorado, as well as their residents, could benefit from more frequent surveys. “***

- Expect surveyors to focus on Infection Control, Life Safety, and Emergency Preparedness.
- Expect stricter enforcement and potentially more severe penalties for violations in these areas.
- I suspect that based on this OIG report, CDPHE will be pressured to FULLY enforce the CMS guidelines related to surveys/inspections and MAY prioritize who and how often to survey based on previous survey results (Risk-based strategy).
- CDPHE survey teams will likely increase the detail of ALL aspects of their investigations and inspections.

[Colorado Could Better Ensure That Nursing Homes Comply With Federal Requirements for Life Safety, Emergency Preparedness, and Infection Control](#)



## February Medical Director Report

[hhs.gov](https://www.hhs.gov)



## Report in Brief

Date: February 2024  
Report No. A-07-22-07009

U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES  
**OFFICE OF INSPECTOR GENERAL**



### Why OIG Did This Audit

In 2016, the Centers for Medicare & Medicaid Services (CMS) updated its life safety and emergency preparedness regulations related to health care facilities to improve protections for all individuals enrolled in Medicare and Medicaid, including those residing in long-term care facilities (nursing homes). The updates expanded requirements related to sprinkler systems, smoke detector coverage, and emergency preparedness plans. Additionally, facilities were required to implement an infection control program.

Our objective was to determine whether Colorado ensured that selected nursing homes in Colorado that participated in the Medicare or Medicaid programs complied with Federal requirements for life safety, emergency preparedness, and infection control.

### How OIG Did This Audit

Of the 219 nursing homes in Colorado that participated in the Medicare and Medicaid programs, we selected a nonstatistical sample of 20 nursing homes for our audit based on location and certain risk factors, including multiple high-risk deficiencies that Colorado reported to CMS.

We conducted unannounced site visits at the 20 nursing homes from September through November 2022. During the site visits, we checked for life safety, emergency preparedness, and infection control deficiencies.

## Colorado Could Better Ensure That Nursing Homes Comply With Federal Requirements for Life Safety, Emergency Preparedness, and Infection Control

### What OIG Found

Colorado could better ensure that nursing homes in Colorado that participated in the Medicare or Medicaid programs comply with Federal requirements for life safety, emergency preparedness, and infection control. During our onsite visits, we identified deficiencies related to life safety, emergency preparedness, or infection control at all 20 nursing homes that we audited, totaling 556 deficiencies. Specifically, we identified 165 deficiencies related to life safety requirements, 210 deficiencies related to emergency preparedness requirements, and 181 deficiencies related to infection control requirements. As a result, the health and safety of residents, staff, and visitors at the 20 nursing homes are at an increased risk during a fire or other emergency, or in the event of an infectious disease outbreak.

The identified deficiencies occurred because of inadequate oversight by Colorado and by nursing home management, frequent management and staff turnover at the nursing homes, inadequate oversight by the State survey agency, and frequent State survey agency staff turnover. In addition, the State survey agency had limited resources to conduct surveys of all nursing homes, including those with a history of multiple high-risk deficiencies, more frequently than was required by CMS. Finally, although not required by CMS, Colorado does not require relevant nursing home staff to participate in standardized life safety training programs despite CMS having a publicly accessible online learning portal with appropriate content.

### What OIG Recommends and Colorado Comments

We recommend that Colorado follow up with the 20 nursing homes reviewed in this audit to ensure that corrective actions have been taken regarding the life safety, emergency preparedness, and infection control deficiencies we identified; work with CMS to develop a risk-based approach to identify nursing homes at which surveys would be conducted more frequently, such as those with a history of multiple high-risk deficiencies or frequent management turnover; and work with CMS to develop standardized life safety training for nursing home staff.

Colorado concurred with our first recommendation and described corrective actions it would take. Colorado did not concur with our second recommendation and neither concurred nor nonconcurred with our third recommendation. We maintain that our findings and all our recommendations are valid.



[OIG-reportDownload](#)

## Keep Calm and Carry On: Managing Residents' Anxiety While Managing Our Own

### Let's Consider a Scenario

***A male resident has been observed to be agitated, short-tempered, and angry. He has been pacing the rehabilitation hall, tends to raise his voice at others, and is often seen watching the exit door. He has been choosing to not take his medications again for the third day in a row, and you observe an interaction between him and the medication aide.***

***The resident is sitting in the corner of his room, and the medication aide is insisting that he takes his medications. He begins to yell at her, calling her names and telling her to leave. She raises her voice to say, "I'm not leaving until you take your medications." The resident responds by rising from his chair, seizing the medications, and throwing them across the room. The medication aide exits the room and tells her coworkers that she refuses to work with the resident again.***

***The resident is referred to a psychologist. His initial evaluation determined that he had smoked two packs of cigarettes per day before his admission, and he has been without nicotine for almost a week. He also is exhibiting moderate to severe anxiety symptoms and symptoms of PTSD related to combat exposure during the Korean War. He reports that he had been taking anxiety medication at home but has not been receiving it.***



***The medication aide in this scenario walked into the situation likely vulnerable to being triggered. She has a personal history of verbal abuse from her significant other and is frequently yelled at in her home life. She stayed to work an extra shift this day because the facility was short-staffed. She was also feeling pressure to hand out the medications quickly so she could leave soon to pick up her kids from school.***

***If we step back from this scenario, we can see that both the resident and the staff member brought predisposing factors into the interaction, which made both more susceptible to respond in an emotional manner at a stressful moment. Had the medication aide recognized that the resident's agitation and anger could be anxiety related – that he was likely experiencing nicotine withdrawal, and that he has PTSD symptoms and may have felt cornered with his back against the wall – she could have used alternative responses. The medication aide, who feels stressed and emotionally triggered when yelled at, also could have politely stepped out and asked a coworker to assist in this potentially triggering situation.***

***[Keep Calm and Carry On: Managing Residents' Anxiety While Managing Our Own - Caring for the Ages](#)***

• **Precipitating Events:**

- Recent major medical diagnosis/terminal illness, surgery, or accident
- Significant change in medical status
- A new environment with strangers
- History of substance use
- Comorbid medical conditions



- Exposure to traumatic events

• **Symptoms:**

- Agitation, nervousness, irritability, or impatience
- A short temper
- Rapid breathing
- Verbalizing worrisome thoughts
- Restlessness
- Not wanting to leave their room
- Refusing therapies
- Avoiding interaction with others
- Difficulty concentrating
- Exaggerated startle response
- Insomnia
- Repeatedly asking to smoke
- Yelling/screaming
- Extreme focus on somatic issues
- Unprovoked physical aggression

**Interventions:**

- **CALM:** Your calmness can promote others to be calm as well.
- **Music:** For residents with dementia, research supports the use of *playing music* (*Gerontologist* 2019;59:e731-e742) to assist with managing anxiety symptoms. Music can lower heart rate, blood pressure, and other autonomic responses. Playing calming music could be beneficial for staff as well.
- **Box Breathing:** If Navy SEALs use it, it must work. Extending your exhalation triggers the parasympathetic nervous system which may lower heart rate and sympathetic drive.



- Inhale to a 4 count, hold breath for 4 seconds, exhale to a 6-10 count, hold for 4 seconds....repeat.
  - **Trigger:** Is there a modifiable trigger to change or avoid?
  - **Exercise:** Physical activity can help balance mood and calm the nervous system.
  - **Generalized Anxiety Disorder 7-item scale (GAD-7)** — [Printable copy of GAD 7](#)
  - **PTSD Screen (PC-PTSD-5)** — [The Primary Care PTSD Screen for DSM-5 \(PC-PTSD-5\) \(va.gov\)](#)
  - **Counseling**
  - **Medications to treat Anxiety:** First line - SSRI and SNRI.
  - **Psychiatry Evaluation**
- 

## Mini Nutritional Assessment and Albumin

[How to Try This: The Mini Nutritional Assessment : AJN The American Journal of Nursing \(lww.com\)](#)

I have seen facilities utilize this Mini Nutritional Assessment as part of the admissions process. I found it a useful review and a tool that I refer to occasionally.

I would also remind you that the use of serum albumin for assessing nutritional status is not sensitive or specific as albumin can be affected by many disease states. It is a negative acute phase reactant.... inflammation (from many causes) reduces albumin/prealbumin.

- Many factors affect the albumin and prealbumin level.
- Normal prealbumin has been seen in studies of patients with BMIs as low as



12.9.

- Prealbumin is not always responsive to nutritional interventions.
- Changes in albumin/prealbumin do not always correlate with outcomes.

[pre-albumin is not useful to assess nutrition.](#)

[how\\_to\\_try\\_this\\_the\\_mini\\_nutritional\\_assessment.30-1Download](#)

## **Summary Recommendation: Do NOT use Urine PCR testing. Stick to CBC, UA with C&S.**

***I have serious concerns that Urine PCR testing will lead to increased inappropriate use of antibiotics. If the physician/NP/PA does not understand the intricacies of the test, it is not unreasonable to expect that antibiotics may be prescribed when they are not needed, the wrong antibiotic may be prescribed, or an antibiotic with a broader than needed spectrum may be used. All of these cases increase the potential for antibiotic resistance and unnecessary antibiotic prescribing.***

***In general, PCR testing is VERY sensitive. It will pick up ANY bacteria that is in the sample. In urine, it will amplify contaminants, colonizers, as well as pathogenic organisms.***

***Urine PCR tests do NOT provide organism antibiotic***



***sensitivities. The test identifies bacterial genes known to cause antibiotic resistance, but it does NOT provide clinically useful antibiotic sensitivities. The existence of a drug resistant gene in one single non-pathologic, contaminating organism could lead one to believe that a powerful broad-spectrum antibiotic is needed when in fact it is not.***

***The report will usually provide antibiotic recommendations based on genetic analysis (not sensitivities). I suspect that these recommendations are primarily aimed at ensuring that there is no liability taken on by the testing company. Thus, the recommendations are often for broader spectrum antibiotics than may be clinically necessary in order to limit medico-legal liability.***

***CDPHE did a pretty good presentation on Urine PCR Testing....see below.***

***[Urine PCR Testing - docShepherd](#)***

---

## Fun Facts:

### ***Hyperkatifeia***

the hypersensitive, negative emotional state associated with withdrawal from a drug. Opioid induced hyperalgesia might be an example. Increased anxiety during benzo withdrawal is another example.

***Hyperkatifeia (derived from the Greek katifeia for dejection or negative emotional state) is defined as an increase in***



***intensity of the constellation of negative emotional or motivational signs and symptoms of withdrawal from drugs of abuse. This overactive negative emotional state is also hypothesized to sensitize with repeated drug exposure and withdrawal and drive an increase in the role of negative reinforcement in maintaining substance use as addiction develops.***

**[hyperkatifeia Optimized \(youtube.com\)](#)**

[Addiction as a Coping Response: Hyperkatifeia, Deaths of Despair, and COVID-19 | American Journal of Psychiatry \(psychiatryonline.org\)](#)

## References

- [Colorado Could Better Ensure That Nursing Homes Comply With Federal Requirements for Life Safety, Emergency Preparedness, and Infection Control \(hhs.gov\)](#)
- [How to Try This: The Mini Nutritional Assessment : AJN The American Journal of Nursing \(lww.com\)](#)
- [Stewardship strategies for interpretation of novel urinary diagnostics \(docshepherd.com\)](#)
- [Urine Polymerase Chain Reaction \(PCR\) Based Testing Guidance Document \(wa.gov\)](#)
- [Brief structured respiration practices enhance mood and reduce physiological arousal - PMC \(nih.gov\)](#)
- [A Brief Measure for Assessing Generalized Anxiety Disorder: The GAD-7 | Anxiety Disorders | JAMA Internal Medicine | JAMA Network](#)
- [Psychometric Properties of the General Anxiety Disorder 7-Item \(GAD-7\) Scale](#)



[in a Heterogeneous Psychiatric Sample - PMC \(nih.gov\)](#)

- [Addiction as a Coping Response: Hyperkatifeia, Deaths of Despair, and COVID-19 | American Journal of Psychiatry \(psychiatryonline.org\)](#)
- [Drug Addiction: Hyperkatifeia/Negative Reinforcement as a Framework for Medications Development - PMC \(nih.gov\)](#)
- [hyperkatifeia Optimized \(youtube.com\)](#)
- [Transition to drug addiction: a negative reinforcement model based on an allostatic decrease in reward function | Psychopharmacology \(springer.com\)](#)